



A Review of the Role of Fungi in Wood Decay of Forest Ecosystems

Bruce G. Marcot¹

Abstract

Fungi are key players in the health, diversity, and productivity of forest ecosystems in Pacific Northwest forests, as mycorrhizal associations, pathogens, decomposers, nontimber resources, and food resources for wildlife. A number of invertebrate species are associated with wood decay fungi, serve as vectors for fungal pathogens, or are fungivorous (consume fungi) and influence rates of wood decay and nutrient mineralization. In Washington and Oregon, 31 wildlife species among 8 families are fungivores, and at least 14 wildlife species disperse fungi. Down wood can provide nurse substrates for seedlings and beneficial mycorrhizal fungi, refuges from pathogenic soil fungi, sources of nutrients for decay fungi, and substrates supporting overall fungal diversity. Presence, density, distribution, and diversity of fungi are influenced by forest stand management practices, forest age class, and effects of fire. Old forests provide for a suite of rare fungi species. Old legacy trees retained during forest harvest can provide some degree of conservation of beneficial and rare fungi. Fungi can be difficult to detect and monitor; surveying for fungi at various times of the year, for multiple (at least 5) years, and by including hypogeous (belowground) samples, can improve detection rates. Studies are needed in the Pacific Northwest to quantify the amount of down wood—number of pieces, sizes, total biomass, percentage of forest floor cover, and other attributes—necessary for maintaining or restoring fungal biodiversity and viable levels of individual fungi species, especially rare species.

Keywords: fungi, mushrooms, mycorrhizae, down wood, coarse woody debris, wood decay, nontimber forest product, fungivores, old forests, monitoring, fire effects.

¹ **Bruce G. Marcot** is a research wildlife biologist, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, 620 SW Main St., Suite 400, Portland OR 97208, bmarcot@fs.fed.us.



Introduction

This report summarizes research and knowledge of the role of fungi in wood decay in the forest ecosystems of Washington and Oregon, and serves to complement the website “DecAID, the Decayed Wood Advisor for Managing Snags, Partially Dead Trees, and Down Wood for Biodiversity in Forests of Washington and Oregon.”² DecAID is a synthesis of data and research results and a planning tool to help inform the management of wood decay elements for biodiversity and species conservation. DecAID describes, in part, fungi associated with decayed wood in Washington and Oregon, including a summary of their ecological roles, the importance of dead wood to fungi, and considerations for the maintenance of fungal biodiversity.

What Are Fungi?

Formally, the term *fungi* as used here refers to the general taxonomic group of organisms that includes rusts, smuts, mildews, molds, yeasts, and mushrooms, and my focus in this review is largely on the mushrooms associated with wood decay. Fungi most associated with wood decay are the filamentous species of Basidiomycota and Ascomycota (Arnstadt et al. 2016, Swift 1982).

More casually, **fungi** also can include the fungus-like slime molds and water molds. Although not discussed here, these nonetheless can be important ecologically and economically, and are more often considered in forest management under pathogen and disease programs. For example, the water mold *Phytophthora ramorum* is responsible for sudden oak death, a forest disease causing widespread killing of oaks and other trees in the Pacific Northwest (Cobb et al. 2012, Rizzo and Garbelotto 2003).

This review covers the various roles and relationships of fungi in wood decay in forests of the Pacific Northwest region of the United States. I also include references to studies conducted outside the Pacific Northwest when local research on specific topics is unavailable.

Ecological Functions of Fungi

Fungi play a number of ecological roles in forest ecosystems that affect the health, diversity, productivity, and development of their biotic communities. Such roles include mycorrhizal associations with vascular plants, pathogens of commercial tree species, decomposers of coarse organic material, and food resources for wildlife.

² <https://www.fs.fed.us/r6/nr/wildlife/decaid/>.

Mycorrhizal Associations

Mycorrhizal fungi consist of strings of hyphae that form mutualistic symbiotic relationships with roots of vascular plants, including trees of commercial value, and that aid the plant in nutrient and water uptake, while the fungi benefit by receiving carbon. Two forms of mycorrhizae are those that grow hyphae from a mantle surrounding the plant roots (ectomycorrhizae) and those with mycelia that embed within the root tissue itself (endomycorrhizae). Allen (1991), O'Dell et al. (1993), and Smith and Read (1997) provided reviews of the structure and function of mycorrhizal fungi.

Fungi As Pathogens

Fungi can also act as pathogens on trees, serving as a cause of tree mortality and altering forest stand structure by opening canopy gaps that, in turn, allow sunlight to penetrate to the forest floor, spurring growth of understory plants and increasing or altering the diversity of plant species (Holah et al. 1993) and other fungi (Christensen 1989). Pathogenic fungi contribute to the accumulation of dead and decaying wood in a forest.

Fungi can create decay in living trees that may be exploited as habitat by some animals. For example, pathogenic fungi such as heartrot fungi can create habitat conditions for primary and secondary cavity-nesting wildlife species and can alter nutrient cycling (Hennon 1995).

Fungi As Decomposers

Fungi associated with down wood are saprobic, meaning that they derive nutrients from decaying organic material. One such species in the Pacific Northwest is orange jelly (*Dacrymyces chrysospermus*), found on decaying logs of Douglas-fir (*Pseudotsuga menzeisii* (Mirb.) Franco) (fig. 1). Other unique fungi associated with down wood and wood decay in the Pacific Northwest are the bird's nest fungus, *Nidula niveotomentosa* (fig. 2), and the veined cup, *Disciotis venosa* (fig. 3).

Fungi found in decaying wood, litter, and duff serve to recycle nutrients (Fogel and Hunt 1983, Hattenschwiler et al. 2005), particularly nitrogen and carbon, as well as minerals, which can then be used by other organisms. Such decomposition processes also serve to physically and chemically break down soil organic matter and alter soil structure. In coarse down wood, wood fungi help mobilize nitrogen, phosphorus, and potassium during the early decay stages (Harmon et al. 1994). Wood decomposition in German forests of European beech (*Fagus sylvatica* L.),



Bruce G. Marcot

Figure 1—Orange jelly mushroom, *Dacrymyces chrysospermus* (previously *D. palmatus*), found on a down log of Douglas-fir in the Cascade Mountains of southwestern Washington.



Bruce G. Marcot

Figure 2—Bird’s nest fungus, *Nidula niveotomentosa*, on a moist Douglas-fir log in the central coast range of Oregon. This unique fungal structure consists of a nest cup called a peridium, that holds “egg” structures called peridioles which contain spore bodies called gleba. In bird’s nest fungi, the peridioles are held in place in the cups with a gelatinous glue-like material until they disperse from splashing raindrops. Species of *Nidula* can reproduce both sexually and asexually, and they produce a ketone chemical with the flavor of raspberry.

Norway spruce (*Picea abies* (L.) Karst.), and Scots pine (*Pinus sylvestris* L.) is dominated by white-rot fungi (*Phanerochaete chrysosporium*), which breaks down lignin in wood (Arnstadt et al. 2016).

In cool temperate and subalpine forests of Japan, Osono (2015) found that litter decomposition was more affected by the presence of specific fungal families than by the type of litter. Fungi of Basidiomycetes had higher rates of lignin breakdown than did fungi of Xylariaceae.



Figure 3—Veined cup fungus, possibly *Disciotis venosa*, in forest litter and down wood fragments, in a Douglas-fir forest of the southern Washington Cascade Mountains. This is one of several brown-colored cup fungi. Although related to the much-sought morel mushrooms, veined cups are likely toxic if eaten raw. They are partially mycorrhizal and thus can play a role in maintaining tree productivity and forest health.

In western Montana, Harvey et al. (1981) found that soil organic matter ≤ 45 percent by volume of the top 30 cm of soil was associated with increased numbers of ectomycorrhizae, but at >45 percent the numbers decreased, and the relationship of soil organic matter and ectomycorrhizae was more salient in dry rather than moist sites.

In studying the role of fungi in decomposition of oak stumps, van der Wal et al. (2015) reported finding unique fungal communities in freshly cut trees and in younger stumps, and noted that old stumps harbored more random assortments of fungal species. They also found that ascomycete fungi likely play a prominent role in wood decay, but stated that further testing is needed, and that better understanding of the fungal roles in wood decay can help improve estimates of carbon sequestration of forests. In southern Sweden, Tyler (1992) likewise found distinct communities of ectomycorrhizal fungi associated with early decay stages of hardwoods.

Fungi As Nontimber Forest Products

Many fungi—particularly aboveground fruiting mushrooms such as chanterelles, morels, matsutake, boletes, truffles, ganoderma (reishi), and others—are sought as food sources, for medicinal use, and by recreational collectors in an expanding

industry (Amaranthus and Pilz 1996, Molina et al. 1993, Pilz et al. 1998, Schlosser and Blatner 1995). Kucuker and Baskent (2017) developed a simulation-based decision-support model to assess the effects of forest management intensity on mushroom occurrence and production. Although developed for northwest Turkey, their system may hold potential for guiding multiple-use forest management in the Pacific Northwest. In a Scots pine forest of central Spain, intensive collection of seasonal sporocarps (aboveground fruiting bodies) of king boletes (*Boletus edulis*) during four productive seasons did not significantly reduce its belowground mycelium biomass, so that the mushroom was able to sustain its productivity (Parladé et al. 2017). This may have implications for monitoring, discussed further below.

Fungi As Food Resources for Wildlife

Fungi themselves are ingested by a wide variety of invertebrate and vertebrate wildlife (Fogel and Trappe 1978, Ingham and Molina 1991, Maser et al. 1978), as discussed more fully in the next section.

Fungi and Invertebrates

Furniss and Carolin (1980) provided a number of examples of insect associations with fungi in forests of the Western United States, as follows. Bark beetles are associated with trees weakened or killed by root-rotting fungi such as Porioa root rot (*Phellinus weirii*), annosus root rot (*Fomes annosus*), and shoestring rot (*Armillaria mellea* and *Phytophthora lateralis*). Some insects, including the smaller European elm bark beetle (*Scolytus multistriatus*), disperse disease-causing fungi, thereby infecting healthy trees. Stain fungi are introduced into weakened trees by bark beetles (especially the western balsam bark beetle, *Dryocoetes confusus*); ambrosia beetles (subfamilies Scolytinae and Platypodinae of Curculionidae); and wood borers (many species and families), the last of which can also mine in sound wood and thereby increase the penetration of wood-rotting fungi in down trees and logs. Ambrosia beetles in particular disperse, introduce, and feed on ambrosia fungi (*Ambrosiella* and *Raffaelea*) and can be highly fungi species-specific. Fir engraver beetles (*Scolytus* spp.) can disperse and introduce brown-stain fungus (*Trichosporium symbioticum*). Some bark beetles (*Gnathotrichus sulcatus*) store and disseminate the symbiotic fungi *Ambrosiella sulcati* and *Raffaelea sulcati*, and the larvae of some horntail insects (*Sirex* and *Urocerus*) feed upon the symbiotic fungi *Amylostereum*. Subterranean termites that comminute (chew) wood fiber are attracted to the wood-decaying fungus *Lenzites trabea*. Among invertebrates associated with yeasts are roundheaded beetles (*Dendroctonus* spp.), bark beetles, and carpenter ants (*Camponotus* spp.). Silver fir beetles (*Pseudohylesinus sericeus*)

can be commonly associated with brown-stain fungi and root-rotting fungi, including *Armillaria mellea*, *Fomes annosus*, and *Phellinus weirii*.

In general, wood-boring insects are known to transport many fungal genera (Schowalter 2000). Ulyshen (2016) reported that invertebrates that are particularly influential in promoting wood decomposition include wood-boring beetles (Coleoptera) and termites (Termitoidae), especially fungus-farming macrotermitines. In a broad study of 13 temperate tree species, Kahl et al. (2017) found that wood decay rates were mediated by enzyme activity and diversity of beetle species. Wood decays more rapidly when it incurs decay fungi introduced by wood-boring beetles, wasps, and termites than when it is initially inoculated with mold fungi by bark and ambrosia beetles (Schowalter 2000). This is because mold fungi can catabolize carbohydrates and thereby inhibit later colonization of decay fungi.

Species interactions that affect changes in fungal and insect communities during wood decay are, in general, poorly understood, and long-term studies are needed. In a boreal forest in central Sweden, Weslien et al. (2011) found that a bark beetle (*Hylurgops palliatus*) and a wood-borer (*Monochamus sutor*) colonized stumps during the first year following cutting; their saproxylic (decaying or dead wood-dependent) functions were mediated by the wood-decaying fungus *Fomitopsis pinicola*, which eventually provided habitat in the stumps 10 years later for a rare, wood-living beetle, *Peltis grossa*. Thus, the researchers suggested this as an example of managing for rare or threatened insect species by understanding the links between saproxylic taxa such as the beetles and the fungi.

Some members of the darkling beetle family Tenebrionidae are associated with fungi (White 1983) (fig. 4). For example, the forked fungus beetle *Bolitotherus cornutus* is nocturnal; during the day it inhabits hard shelf fungi or crevices where the fungi are attached. The darkling beetles *Diaperis* spp. and *Playtdema* spp. occur under bark and in fungi. The aptly named handsome fungus beetles of family Endomychidae, such as the Idaho handsome fungus beetle *Mycetina idahoensis*, occur under bark in rotting wood and in fungi on which they feed (Haggard and Haggard 2006).

Fungivorous insects are typically associated with late-successional forests (Schowalter 2000) and can influence the diversity of fungi in decaying wood in both managed and natural forests (Muller et al. 2002). Fungivorous springtails apparently serve to transfer secondary metabolites (catalpol, an iridoid glucoside) from host plants to arbuscular endomycorrhizal fungi (Duhamel et al. 2013). This functions in the fungi to prevent it from being grazed. In this triad of relationships, the springtails benefit from the fungal food source, the fungi benefits from avoiding grazing, and the host plant benefits from using the symbiotic fungi to absorb soil nutrients.



© Joyce Gross, used by permission

Figure 4—A horned fungus beetle (*Bolitotherus cornutus*) (family Tenebrionidae, darkling beetles), found in the woods of eastern North America. This species feeds on the tissue of bracket fungi (polypores) on dead or dying tree trunks. It lays its eggs within the fungi, and the larvae pupate within the bracket or in nearby soil. Other fungivorous species of this family, such as the broad-necked darkling beetle (*Coelocnemis californicus*, previously *C. dilaticollis*), are found in the Northwest but are poorly studied.

In other symbiotic relationships, Macrotermitinae termites deposit all their feces in their tended gardens of the fungus *Termitomyces* spp. (Basidiomycetes). Individual termite species of this group tend to be associated with, and feed only on, specific species of these fungi (Edwards 2000).

Nutrients in woodland soils can be greatly affected by some invertebrate associations with fungi, as reported by Crowther et al. (2011a). Invertebrate grazers in soil can determine the composition of fungal decomposer communities. For example, isopods were found to feed selectively on the cord-forming fungus *Resinicium bicolor*, thus preventing the competitive exclusion of two fungi species in soil and wood. Similar mediating functions were also observed with soil nematodes. Thus, conditions affecting soil invertebrates can also affect their influence on fungal communities and associated nutrient cycles. Also, invertebrate fungivory can influence decay rates of wood and nutrient mineralization and decomposition (Crowther et al. 2011b).

Some mycorrhizal fungi produce non-nitrogenous chemical defenses, including pyrethroids that are toxins absorbed through insect exoskeletons (Schowalter 2000).

Fungi and Wildlife

In Washington and Oregon, some 31 wildlife species among 8 families (all mammals) are known to be fungivores (table 1).

Table 1—Fungivorous wildlife species of Washington and Oregon

Family	Common name	Scientific name
Cervidae	Black-tailed deer	<i>Odocoileus hemionus columbianus</i>
Cervidae	Mule deer	<i>Odocoileus hemionus hemionus</i>
Cervidae	Rocky Mountain elk ^a	<i>Cervus elaphus nelsoni</i>
Cervidae	Roosevelt elk ^a	<i>Cervus elaphus roosevelti</i>
Dipodidae	Pacific jumping mouse	<i>Zapus trinotatus</i>
Geomyidae	Camas pocket gopher	<i>Thomomys bulbivorus</i>
Geomyidae	Northern pocket gopher	<i>Thomomys talpoides</i>
Geomyidae	Townsend's pocket gopher	<i>Thomomys townsendii</i>
Muridae	Bushy-tailed woodrat	<i>Neotoma cinerea</i>
Muridae	Canyon mouse	<i>Peromyscus crinitus</i>
Muridae	Columbian mouse ^a	<i>Peromyscus keeni</i>
Muridae	Creeping vole	<i>Microtus oregoni</i>
Muridae	Deer mouse ^a	<i>Peromyscus maniculatus</i>
Muridae	Pinon mouse	<i>Peromyscus truei</i>
Muridae	Southern red-backed vole ^a	<i>Myodes gapperi</i>
Muridae	Western red-backed vole ^a	<i>Myodes californicus</i>
Muridae	White-footed vole ^a	<i>Arborimus albipes</i>
Ochotonidae	American pika	<i>Ochotona princeps</i>
Sciuridae	Douglas' squirrel ^a	<i>Tamiasciurus douglasii</i>
Sciuridae	Golden-mantled ground squirrel	<i>Spermophilus lateralis</i>
Sciuridae	Least chipmunk ^a	<i>Tamias minimus</i>
Sciuridae	Northern flying squirrel ^a	<i>Glaucomys sabrinus</i>
Sciuridae	Red squirrel ^a	<i>Tamiasciurus hudsonicus</i>
Sciuridae	Townsend's chipmunk	<i>Tamias townsendii</i>
Sciuridae	Siskiyou chipmunk ^a	<i>Tamias siskiyou</i>
Sciuridae	Western gray squirrel ^a	<i>Sciurus griseus</i>
Sciuridae	Yellow-pine chipmunk	<i>Tamias amoenus</i>
Soricidae	Pacific shrew	<i>Sorex pacificus</i>
Soricidae	Trowbridge's shrew	<i>Sorex trowbridgii</i>
Soricidae	Vagrant shrew	<i>Sorex vagrans</i>
Suidae	Feral pig	<i>Sus scrofa</i>

^a Also disperses fungi.

Source: Jacobs and Luoma (2008), O'Neill et al. (2001).

Note: taxonomy is from Hayssen, V.; Mammalian Species, American Society of Mammalogists, <http://www.science.smith.edu/departments/Biology/VHAYSEN/msi/default.html>.

Some fungi are dispersed on the beaks of foraging and cavity-excavating woodpeckers (Jusino et al. 2016), thereby serving to inoculate live and dead trees. Fungi such as truffles and their ectomycorrhizal sporocarps are key food resources for northern flying squirrels (*Glaucomys sabrinus*) (Lehmkuhl et al. 2004); in turn, flying squirrels are a key prey species of northern spotted owls (*Strix occidentalis caurina*) in parts of the owl's range.

Some fungi are highly detrimental to some species of wildlife, such as the deadly amphibian disease of chytridiomycosis caused by the fungus *Batrachochytrium dendrobatidis*, and white-nose syndrome, which is debilitating and deadly to bats, caused by the fungus *Pseudogymnoascus destructans*. However, there is no evidence that these fungal pathogens are related to wood decay.

Fungi-dispersing wildlife in this region (table 1) number at least 14 species, including American pika (*Ochotona princeps*). Species of deer and elk can disperse fungi through their pellets (fig. 5). Small mammals, such as white-footed voles (Manning et al. 2003), are among the species that are documented as dispersers of mycorrhizal fungi (Luoma et al. 2003, Maser et al. 1978).

In general, fungi species with hypogeous sporocarps (which release spores below ground), such as truffles, depend on animals for dispersal. Jacobs and Luoma (2008) studied four forest rodents (Townsend's chipmunk, Siskiyou chipmunk, western red-backed vole, and southern red-backed vole) that serve both as dispersers of truffles, including *Rhizopogon*, and as prey for northern spotted owls. They found that isolated green-tree retention in harvest blocks reduced consumption of truffles by the voles, and that the impact could be offset by including green-tree aggregates within a dispersed retention matrix.

Maser and Maser (1988) reported that all squirrels of 5 genera and 8 species in Oregon conifer forests are mycophagous (eat fungi), particularly consuming belowground fruiting bodies of at least 26 genera of mycorrhizal fungi. Northern flying squirrels proved to be a nearly obligate fungivore. In general, they found that squirrels may be vital links involving belowground mycorrhizal fungi, nitrogen-fixing bacteria, yeast, and conifer trees.

Marcot (2002) demonstrated how a "functional web" can be depicted for wildlife associated with various wood decay elements (snags, down wood, litter, duff, mistletoe brooms, dead parts of live trees, hollow living trees, natural tree cavities, bark crevices, and live remnant or legacy trees), including wildlife species responsible for dispersing fungi, in Washington and Oregon.



Bruce G. Marcot



Bruce G. Marcot

Figure 5—Fungi dispersed via pellet droppings from Rocky Mountain elk. Tower and Summit Burn, Malheur National Forest, eastern Oregon.

Wood Decay and Fungi

The dynamics of wood decay are linked closely to the presence and ecological functions of fungi. Decay of snags and down wood proceeds through a series of stages marked by degree of wood breakdown, changes in the diversity of associated biota, progressions of nutrient transformations, and other processes. Spies and Cline (1988) and Maser et al. (1979) provided a 5-category classification system of wood

decay in down wood, progressing from recently downed wood with intact bark, branches, and twigs (decay class I) to advanced states of wood breakdown into soft textures of duff (decay class V).

Throughout this mini-successional sequence of wood decay, fungi, along with mesoarthropods and other species, play key physical and biochemical roles in wood decomposition and nutrient cycling. In particular, in young and old Douglas-fir stands, the ectomycorrhizal fungus *Piloderma fallax* increases in occurrence in relation to percentage of cover of down wood of the advanced decay class V. The presence of truffle and false truffle fungi has also been shown to be associated with proximity to (within 1 m of) down wood (Amaranthus et al. 1994).

Down wood, throughout its decay sequence, also serves to retain moisture, which promotes growth of ectomycorrhizae (Amaranthus et al. 1989; Harmon and Sexton 1995; Harvey et al. 1976, 1978), and which thereby serves as refugia for seedlings and mycorrhizal fungi. Such “reservoir” functions of down wood can be particularly salient in xeric forests and during dry seasons, providing for establishment of beneficial mycorrhizal fungi as a forest stand regrows (O’Hanlon-Manners and Kotanen 2004) and serving as “nurse logs” for seedlings of vascular plants (Harmon and Franklin 1989, Kropp 1982) such as western hemlock (*Tsuga heterophylla* (Raf.) Sarg.), Engelmann spruce (*Picea engelmannii* Parry ex Engelm.), and subalpine fir (*Abies lasiocarpa* (Hook.) Nutt.) (Brang et al. 2003). Nurse logs also can act as refuges from pathogenic soil fungi (O’Hanlon-Manners and Kotanen 2004).

Decaying down wood provides nutrients for decay fungi and pathogens. Studies in North America and Scandinavia both reveal that high diversity of wood-decay fungal species is associated with the presence of large down wood (Bader et al. 1995, Crites and Dale 1998, Høiland and Bendiksen 1996, Kruys et al. 1999, Ohlson et al. 1997, Wästerlund and Ingelög 1981). Høiland and Bendiksen (1996) found that rare wood-inhabiting fungal species occurred primarily on long (mean 11 m) and well-decayed (average decay class III) down wood. Kruys and Jonsson (1999) found that fungal species diversity is associated with total surface area of down wood.

Fungi in Forest Management

Functioning forest ecosystems in the Pacific Northwest depend on the diversity and viability of fungal species. The presence, density, distribution, and diversity of fungi are influenced by forest stand management practices and by forest age

class (Cázares et al. 1998, Clarkson and Mills 1994, Colgan et al. 1999, Heithecker and Halpern 2006, O'Dell et al. 1992, Pilz and Perry 1984, Stendell et al. 1999, Trofymow et al. 2003). And, in turn, plant community structures, including wood decay elements, are influenced by fungi in complex feedback systems (van der Heijden et al. 1998).

In a study in France, Paillet et al. (2017) reported that snags, more so than large live trees, provide the bulk of tree microhabitats, including cavities, fungi conks, and bark features, and that strict forest reserves contain a greater abundance of such microhabitats than do managed forests.

Influence of Forest Management Activities

Thinning and clearcutting alter the fungal community and can reduce the production of sporocarps and ectomycorrhizae. Rydin et al. (1997) found that habitat loss and some forest management practices in Europe have led to declines in the diversity of fungi and in the presence of rare fungal species. Berg et al. (1994) reported that many fungal species in Swedish forests are threatened by the loss of old trees and declines in coarse woody debris. Arnstadt et al. (2016) noted that higher intensities of forest management in Germany negatively affect the volume of dead wood and richness of fungal species sporocarps. Parladé et al. (2017) found that clearcutting and partial cutting of Scots pine forests in central Spain equally and sharply reduced the mycelium biomass of king boletes (*Boletus edulis*).

In European Norway spruce stands, Löhmus (2011) studied the influence of clearcutting, planting, and thinning on polypore (bracket) fungi. Results indicated that distinct polypore communities were present in clearcuts but their species richness declined over time and increased again 20 years post-cutting and following tree planting. Most polypore species were found in mature, unmanaged, naturally regenerated stands; thinning reduced species richness by 15 percent; and distinct polypore communities were present in young stands on nutrient-rich soils.

Fungi in Old Forests

Under the Northwest Forest Plan (NWFP) in western Washington and Oregon and northwestern California, the Survey and Manage Program listed 234 rare fungi species found in late-successional and old-growth forests (Molina 2008), many of which are associated with various aspects of wood decay. Molina (2008) noted that some two-thirds of these species also occurred outside late-successional forest reserves under the NWFP, suggesting that conservation of fungal biodiversity may benefit from additional guidelines outside the reserves. More recently, the

Interagency Special Status and Sensitive Species Program of the Pacific Northwest Region of the U.S. Forest Service and Bureau of Land Management has taken over the role and duties of the Survey and Manage Program, including providing an annotated bibliography of rare species of fungi of California, Oregon, and Washington.³

Managing for Fungal Species and Communities

Except for sensitive or listed species, no general guidelines are in place to provide for conservation or restoration of fungal communities, including those associated with wood decay elements. It is known, though, that retention of legacy trees—usually mature or old-growth trees retained during forest harvest operations—can provide some degree of conservation of beneficial fungi such as mycorrhizae (Smith et al. 2000). Retaining green trees has been attributed by Luoma (2001) to retention of the rare truffle *Arcangeliella camphorata*, which is otherwise lost in clearcuts, such as has been demonstrated in southwest Oregon (Amaranthus et al. 1994, Clarkson and Mills 1994). In Washington, Cline et al. (2005) reported that Douglas-fir seedlings nearer (<6 m) to residual mature Douglas-fir trees in recently harvested green-tree retention units had higher species richness and diversity of ectomycorrhizal fungi than did seedlings far from residual trees. They thus suggested that residual mature, legacy trees can maintain or accelerate recovery of ectomycorrhizal fungi following harvest. As well, retained stumps can provide environments for conks and other fungi (fig. 6).

In some cases, active management can help retain or restore desired fungi by deliberately introducing fungi in live trees. This can help foster wood decay and create snags and dead parts of live trees for wildlife habitat, such as demonstrated by Bednarz et al. (2013) and Filip et al. (2011) in forests of Oregon and Washington.



Bruce G. Marcot

Figure 6—Cut stumps, along with coarse and fine down wood and other wood decay elements, can provide substrates for wood-decaying fungi such as these conks of *Fomatopsis pinicola*. Gifford Pinchot National Forest, Washington.

³ <http://www.fs.fed.us/r6/sfpnw/issssp/documents3/cpt-fu-effects-guidelines-att3-annotated-bibliography-2013-10.docx>.

To maintain fungal biodiversity, the habitat and resource associations of multiple species need to be considered. This can be achieved, in part, by providing a range of sizes and decay classes of down wood, although such associations of individual species and their responses to various amounts, patterns, sizes, decay classes, and timing of down wood are poorly known and need much study. In general, though, providing down wood as well as living host plants of the correct ages and species can help maintain fungal diversity.

In Sweden, Edman and Jonsson (2001) and Edman et al. (2007) reported that the spatial distribution of down logs and wood-decaying fungi are influenced by wind and gap-phase dynamics in forests of old-growth Norway spruce. They also found that rare fungi species have specific substrate associations and that temporal variations in the patterns of canopy gaps and down wood abundance can affect fungi biodiversity. White et al. (2012) studied the effect of a massive ice storm in forests of southern Quebec, Canada, which caused forest canopy gap openings that became colonized by wood-rotting fungi, saproxylic insects, salamanders, and other organisms. Such canopy gap dynamics apparently served to maintain the diversity of opening-dependent taxa, including some fungi.

However, in a study of ectomycorrhizal fungi based on epigeous sporocarps in a cedar-hemlock forest of northwest British Columbia, Canada, Durall et al. (1999) found that fungal species richness decreased exponentially as a function of increasing size of forest gap cutblocks, particularly in gaps $>900 \text{ m}^2$. Maximum fungal species richness was found $\leq 7 \text{ m}$ from the forest edge. They suggested sampling both sporocarps and root tips for accurately determining the ectomycorrhizal fungal community.

In a study of northern hardwood forests, Brazeel et al. (2014) found various fungi species associated with a variety of conditions, including stumps, down wood of small ($<20 \text{ cm}$ diameter) through large ($>40 \text{ cm}$ diameter) sizes, well-decayed substrates, minor host tree species, and canopy gaps. In Norway spruce forests of Sweden, Edman et al. (2004) found that fungi was more common in sites rich in down wood, and that fungi species richness was more greatly associated with large logs than with small logs. Crawford et al. (1990) found that filamentous fungi and yeast communities in Douglas-fir logs varied between decay classes III and IV, and that they discovered a total of 18 genera and 36 species of fungi among logs of both decay classes.

Studies are needed in the Pacific Northwest to quantify the amount of down wood—number of pieces, sizes, total biomass, percentage of forest floor cover, and other attributes—necessary for maintaining or restoring fungal biodiversity and viable levels of individual fungi species, especially rare species. Also, fungi tend to

occur in patchy distributions because of the patchy occurrence of their substrates. Providing down wood of various sizes, species, and decay classes in patchy distributions throughout stands in managed forest landscapes may help restore and maintain desired fungal communities.

Surveys of wood-inhabiting fungi in spruce-hardwood forests of central Finland (Juutilainen et al. 2011) found a distinct fungal community in the smallest pieces of down wood; by excluding pieces <1 cm in diameter, fungi species richness, including rare species, was underestimated by 10 percent and occurrences by 46 percent. Surveying fungi only in larger down wood (coarse woody debris) seriously underestimated richness and abundance of dead wood-associated fungi.

It takes time for mycorrhizae to colonize down wood and coarse woody debris, because most mycorrhizal fungi in wood are associated with roots. A good example is *Boletus (Aureoboletus) mirabilis*, which always fruits from decay class IV or V wood, but that is because it is mycorrhizal with the roots of hemlock in the wood. This time delay needed for colonization and association with roots highlights the role and value of retaining some late-seral forests and old legacy trees as refugia and as source material for beneficial fungi (Clarkson and Mills 1994). Otherwise, sources may be relegated to disturbance-resistant fungi spores remaining in soil or in whatever unburned down wood may remain after disturbance (Baar et al. 1999). Still, reappearance of some fungi may appear delayed following disturbance, such as chanterelles (*Cantharellus*) appearing in western hemlock stands after 20 years following disturbance along the Washington coast (Pilz et al. 1998). But once established in appropriate habitat conditions, mycelial colonies of fungi can persist for many years (Dahlberg and Stenlid 1995, De la Bastide et al. 1994, Smith et al. 1992).

Lehmkuhl et al. (2007) discussed a decision-aiding model FuelSolve that can be used to guide management of fuels in forests under multiple objectives such as providing habitat for northern spotted owls and their prey, along with live and dead vegetation, mycorrhizal fungi, and arboreal lichens, as elements of the owl's habitat.

Further ideas on managing Pacific Northwest forests for fungi can be found in Molina et al. (2001).

Monitoring Fungi

Fungi are often difficult to detect, especially for determining the presence of rare, sparsely-distributed, and seldom-fruited species. Most species can be detected only when they produce reproductive structures such as cups, truffles, conks, and mushrooms (fig. 7). Different species may produce such detectible structures at different times and seasons (Luoma 1991), depending on species-specific relationships



Figure 7—Fruiting bodies (sporocarps) of fungi may appear intermittently, seasonally, or rarely, depending on the species, its rarity, and environmental conditions, making monitoring a challenge. (A) Sporocarps of *Galerina marginata*, a most deadly species, on a down Douglas-fir log. (B) Sporocarps from *Mycea mycelia* beneath the log; their mycelia commonly grow from fine woody debris and litter.

to nutrient availability and environmental conditions of temperature, light, pH, and moisture. O'Dell et al. (1996) recommended surveying for fungi at various times of the year, particularly in spring and autumn, for at least 5 years, to provide any assurance of detection.

Lassauce et al. (2011) tested the idea that dead wood volume could be monitored as an index to species richness of saproxylic beetles and fungi in various forest types. However, they found that correlations were only moderately significant and concluded that dead wood volume is likely an imprecise indicator of saproxylic beetle and fungi biodiversity. Further, the efficacy of using dead wood volume to indicate saproxylic beetle diversity differed between boreal and temperate forests, with slightly greater predictability in the former. They suggested that additional landscape-level variables, such as the type and decay class of dead wood, be included in monitoring dead wood and associated organisms. Parladé et al. (2017) suggested that surveys of soil mycelium masses (fig. 8) can usefully indicate the response of some fungi to management activities, and could be useful adjuncts to monitoring sporocarp fruiting bodies of interest to gatherers.



Bruce G. Marcot

Figure 8—Fungi mycelium mass beneath a log. Studies suggest that monitoring just the fruiting bodies (sporocarps, fig. 7) may underestimate fungal community diversity, and that surveying soil mycelium masses can better indicate response of fungi to forest management activities.

Another challenge to monitoring fungi related to wood decay is to identify the appropriate spatial and temporal scales. In a review of studies on saproxylic species (fungi, beetles, and lichens) and associated dead wood distribution in Europe, Sverdrup-Thygeson et al. (2014) identified key information gaps. They found a large variation among taxa of such species in response to spatial and temporal variations in dead wood patterns. They suggested that time-lag effects, in particular, need more study at landscape scales and for listed saproxylic species before firm management guidelines can be developed for them.

Influence of Fire

The main influence of fire on wood decay-associated fungi relates to how much sound or decaying wood is created or destroyed. Prescribed fires and wildfires alike can kill part or all of standing trees which, if not engulfed and fully charred, could provide fungi substrates standing or down. Fire also can eliminate fungi substrates, particularly with piling and burning of forest slash following timber harvests.

In forests of the eastern Cascade Range in Oregon, Smith et al. (2017) studied soil fungal and bacterial communities and biogeochemical processes following severe and less-severe burns. They found that soil fungi and bacteria steadily recolonized following the burns, but with a different community composition between the two fire severities. The greatest difference in fungal and bacterial community composition was evident early after the burns and became more similar over time.

In Swedish forests of Scots pine, Jonsson et al. (1999) compared chronosequences of ectomycorrhizae in stands burned by low-intensity wildfire and unburned late-successional stands. They found most of the common species in all sites, suggesting that ectomycorrhizae exhibit a continuity following low-intensity burning. Importantly, the belowground species composition was not reflected in that of the aboveground sporocarps.

Fungi As a Conservation Challenge

Maintaining and restoring desired wood decay-associated fungi can be quite a challenge for management (O'Dell et al. 1996) given the problems of intermittent detectability, variable dispersal, patchy distributions, and lack of scientific information on species' life histories and habitat requirements. Further challenges include identifying species, the need for taxonomic studies, and incomplete understanding of their ecological functional roles in forest ecosystems. Studies conducted over the past decade have shed light on some fungi species in some geographic areas and forest types of the Pacific Northwest (e.g., see footnote 2).

In a global review of conservation strategies for managing dead wood for biodiversity, Seibold et al. (2015) found many information gaps and, at best, only scattered management guidelines. Their meta-analysis revealed that most studies have focused on early stages of wood decay and that some taxa, including fungi, are underrepresented. The studies do confirm the overall benefits of dead wood for biodiversity, but there is a need for research on advanced decay stages and on the influence on non-saprophytic organisms.

Still, fungi are key players in native and productive forests, and offer important roles in nutrient cycling, food sources, tree production, and maintenance of soil health.

Acknowledgments

This review is based in part on, and is updated from, a previous essay by Tina Dreisbach (2002), posted as part of the DecAID Decayed Wood Advisor version 2.20 (Mellen et al. 2002), and also borrows in part from Marcot et al. (N.d.). The original report by Dreisbach was reviewed by Jane E. Smith, Randy Molina, Thomas O'Dell, Efrén Cázares, and the DecAID Science Team. I acknowledge and appreciate their work. Thanks also to Dan Luoma, Steven Acker, and Kim Mellen-McLean for helpful reviews of the current manuscript.

English Equivalents

When you know:	Multiply by:	To find:
Centimeters (cm)	0.394	Inches
Square meters (m ²)	10.76	Square feet

References

- Allen, M. F. 1991.** The ecology of mycorrhizae. Cambridge, United Kingdom: Cambridge University Press. 184 p.
- Amaranthus, M.P.; Parrish, D.S.; Perry, D.A. 1989.** Decaying logs as moisture reservoirs after drought and wildfire. In: Alexander, E., ed. Proceedings of the Watershed 1989 symposium on stewardship of soil, air, and water resources. R10-MB-77. Juneau, AK: U.S. Department of Agriculture, Forest Service, Alaska Region: 191–194.

- Amaranthus, M.; Trappe, J.M.; Bednar, L.; Arthur, D. 1994.** Hypogeous fungal production in mature Douglas-fir forest fragments and surrounding plantations and its relation to coarse woody debris and animal mycophagy. *Canadian Journal of Botany*. 24: 2157–2165.
- Amaranthus, M.; Pilz, D. 1996.** Productivity and sustainable harvest of wild mushrooms. In: Pilz, D.; Molina, R., eds. *Managing forest ecosystems to conserve fungus diversity and sustain wild mushroom harvests*. Gen. Tech. Rep. PNW-GTR-371. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 42–61.
- Arnstadt, T.; Hoppe, B.; Kahl, T.; Kellner, H.; Krüger, D.; Bauhus, J.; Hofrichter, M. 2016.** Dynamics of fungal community composition, decomposition and resulting deadwood properties in logs of *Fagus sylvatica*, *Picea abies* and *Pinus sylvestris*. *Forest Ecology and Management*. 382: 129–142.
- Baar, J.; Horton, T.R.; Kretz, A.M.; Bruns, T.D. 1999.** Mycorrhizal colonization of *Pinus muricata* from resistant propagules after a stand-replacing wildfire. *New Phytologist*. 143: 409–418.
- Bader, P.; Jansson, S.; Jonsson, B.G. 1995.** Wood-inhabiting fungi and substratum decline in selectively logged boreal spruce forest. *Biological Conservation*. 72: 355–362.
- Bednarz, J.C.; Huss, M.J.; Benson, T.J.; Varland, D.E. 2013.** The efficacy of fungal inoculation of live trees to create wood decay and wildlife-use trees in managed forests of western Washington, USA. *Forest Ecology and Management*. 307: 186–195.
- Berg, A.; Ehnstrom, B.; Gustafsson, L.; Hallingback, T.; Jonsell, M.; Weslien, J. 1994.** Threatened plant, animal, and fungus species in Swedish forests: distribution and habitat associations. *Conservation Biology*. 8(3): 718–731.
- Brang, P.; Moran, J.; Puttonen, P.; Vyse, A. 2003.** Regeneration of *Picea engelmannii* and *Abies lasiocarpa* in high-elevation forests of south-central British Columbia depends on nurse logs. *Forestry Chronicle*. 79(2): 247–252.
- Brazee, N.J.; Lindner, D.L.; D’Amato, A.W.; Fraver, S.; Forrester, J.A.; Mladenoff, D.J. 2014.** Disturbance and diversity of wood-inhabiting fungi: effects of canopy gaps and downed woody debris. *Biodiversity and Conservation*. 23(9): 2155–2172.

- Cázares, E.; Luoma, D.L.; Eberhart, J.; Amaranthus, M.P.; Cray, C.; Dudd, J.; McArthur, M. 1998.** Hypogeous fungal diversity and biomass following salvage logging in Mt. Hood National Forest, Oregon, USA. In: Ahonen-Jonnarth, U., ed. Programme and abstracts of the second international conference of mycorrhiza. Uppsala, Sweden: Swedish University of Agricultural Sciences: 39–40.
- Christensen, M. 1989.** A view of fungal ecology. *Mycologia*. 81: 1–19.
- Clarkson, D.A.; Mills, L.S. 1994.** Hypogeous sporocarps in forest remnants and clearcuts in southwest Oregon. *Northwest Science*. 68(4): 259–265.
- Cline, E.T.; Ammirati, J.E.; Edmonds, R.L. 2005.** Does proximity to mature trees influence ectomycorrhizal fungus communities of Douglas-fir seedlings? *The New Phytologist*. 166(3): 993–1009.
- Cobb, R.C.; Chan, M.N.; Meentemeyer, R.K.; Rizzo, D.M. 2012.** Common factors drive disease and coarse woody debris dynamics in forests impacted by sudden oak death. *Ecosystems*. 15(2): 242–255.
- Colgan, W., III; Carey, A.B.; Trappe, J.M.; Molina, R.; Thysell, D. 1999.** Diversity and productivity of hypogeous fungal sporocarps in a variably thinned Douglas-fir forest. *Canadian Journal of Forest Research*. 29: 1259–1268.
- Crawford, R.H.; Carpenter, S.E.; Harmon, M.E. 1990.** Communities of filamentous fungi and yeast in decomposing logs of *Pseudotsuga menziesii*. *Mycologia*. 82(6): 759–765.
- Crites, S.; Dale, M.R.T. 1998.** Diversity and abundance of bryophytes, lichens, and fungi in relation to woody substrate and successional stage in aspen mixedwood boreal forests. *Canadian Journal of Botany*. 76: 641–651.
- Crowther, T.W.; Boddy, L.; Jones, T.H. 2011a.** Outcomes of fungal interactions are determined by soil invertebrate grazers. *Ecology Letters*. 14(11): 1134–1142.
- Crowther, T.W.; Jones, T.H.; Boddy, L. 2011b.** Species-specific effects of grazing invertebrates on mycelial emergence and growth from woody resources into soil. *Fungal Ecology*. 4: 333–341.
- Dahlberg, A.; Stenlid, J. 1995.** Spatiotemporal patterns in ectomycorrhizal populations. *Canadian Journal of Botany*. 73(Suppl. 1): 1222–1230.
- De la Bastide, P.Y.; Kropp, B.R.; Piche, Y. 1994.** Spatial distribution and temporal persistence of discrete genotypes of the ectomycorrhizal fungus *Laccaria bicolor* (Maire) Orton. *New Phytologist*. 127: 547–556.

- Dreisbach, T. 2002.** Importance of fungi in forest ecosystems. In: DecAID, the Decayed Wood Advisor for managing snags, partially dead trees, and down wood for biodiversity in forests of Washington and Oregon. Version 2.20. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. <https://www.fs.fed.us/r6/nr/wildlife/decaid/pages/Fungi.html>. (May 26, 2017).
- Duhamel, M.; Pel, R.; Ooms, A.; Bücking, H.; Jansa, J.; Ellers, J.; van Straalen, N.M.; Wouda, T.; Vandenkoornhuyse, P.; Kiers, E.T. 2013.** Do fungivores trigger the transfer of protective metabolites from host plants to arbuscular mycorrhizal hyphae? *Ecology*. 94 (9): 2019–2029.
- Durall, D.M.; Jones, M.D.; Wright, E.F.; Kroeger, P.; Coates, K.D. 1999.** Species richness of ectomycorrhizal fungi in cutblocks of different sizes in the interior cedar-hemlock forests of northwestern British Columbia: sporocarps and ectomycorrhizae. *Canadian Journal of Forest Research*. 29: 1322–1332.
- Edman, M.; Jonsson, B.G. 2001.** Spatial pattern of downed logs and wood-decaying fungi in an old-growth *Picea abies* forest. *Journal of Vegetation Science*. 12(5): 609–620.
- Edman, M.; Jönsson, M.; Jonsson, B.G. 2007.** Fungi and wind strongly influence the temporal availability of logs in an old-growth spruce forest. *Ecological Applications*. 17(2): 482–490.
- Edman, M.; Kruys, N.; Jonsson, B.G. 2004.** Local dispersal sources strongly affect colonization patterns of wood-decaying fungi on spruce logs. *Ecological Applications*. 14(3): 893–901.
- Edwards, C.A. 2000.** Soil invertebrate controls and microbial interactions in nutrient and organic matter dynamics in natural and agroecosystems. In: Coleman, D.C.; Hendrix, P., eds. *Invertebrates as webmasters in ecosystems*. Wallingford, Oxon, United Kingdom: CABI Publishing: 141–159.
- Filip, G.; Chadwick, K.; Zambino, P.; Omdal, D.; Ramsey-Kroll, A.; Schmitt, C.; Maffei, H.; Saavedra, A.; Rall, W.; Parks, C. 2011.** Seven- to 14-year effects of artificially inoculating living conifers to promote stem decay and subsequent wildlife use in Oregon and Washington forests. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, Forest Health Protection. 24 p.

- Fogel, R.; Hunt, G. 1983.** Contribution of mycorrhizae and soil fungi to nutrient cycling in a Douglas- fir ecosystem. *Canadian Journal of Forest Research*. 13: 219–232.
- Fogel, R.; Trappe, J.M. 1978.** Fungus consumption (mycophagy) by small animals. *Northwest Science*. 52: 1–30.
- Furniss, R.L.; Carolin, V.M. 1980.** Western forest insects. Misc. Publications 1339. Washington, DC: U.S. Department of Agriculture, Forest Service. 654 p.
- Haggard, P.; Haggard, J. 2006.** Insects of the Pacific Northwest. Portland, OR: Timber Press. 295 p.
- Harmon, M.E.; Franklin, J.F. 1989.** Tree seedlings on logs in *Picea-Tsuga* forests of Oregon and Washington. *Ecology*. 70: 48–59.
- Harmon, M.E.; Sexton, J.; Caldwell, B.A.; Carpenter, S.E. 1994.** Fungal sporocarp mediated losses of Ca, Fe, K, Mg, Mn, N, P, and Zn from conifer logs in the early stages of decomposition. *Canadian Journal of Forest Research*. 24: 1883–1893.
- Harmon, M.E.; Sexton, J. 1995.** Water balance of conifer logs in early stages of decomposition. *Plant and Soil*. 172: 141–152.
- Harvey, A.E.; Jurgensen, M.F.; Larsen, M.J. 1978.** Seasonal distribution of ectomycorrhizae in a mature Douglas-fir/larch forest soil in western Montana. *Forest Science*. 24: 203–208.
- Harvey, A.E.; Jurgensen, M.F.; Larsen, M.J. 1981.** Organic reserves: importance to ectomycorrhizae in forest soils of western Montana. *Forest Science*. 27(3): 442–445.
- Harvey, A.E.; Larsen, M.J.; Jurgensen, M.F. 1976.** Distribution of ectomycorrhizae in a mature Douglas-fir/larch forest soil in western Montana. *Forest Science*. 22: 393–398.
- Hattenschwiler, S.; Tiunov, A.V.; Scheu, S. 2005.** Biodiversity and litter decomposition in terrestrial ecosystems. *Annual Review of Ecology and Systematics*. 36: 191–218.
- Heithecker, T.D.; Halpern, C.B. 2006.** Variation in microclimate associated with dispersed-retention harvests in coniferous forests of western Washington. *Forest Ecology and Management*. 226 (1–3): 60–71.
- Hennon, P.E. 1995.** Are heart rot fungi major factors of disturbance in gap-dynamic forests? *Northwest Science*. 69: 284–292.

- Høiland, K.; Bendiksen, E. 1996.** Biodiversity of wood-inhabiting fungi in a boreal coniferous forest in Sør-Trøndelag County, Central Norway. *Nordic Journal of Botany*. 16: 643–659.
- Holah, J.C.; Wilson, M.V.; Hansen, E.M. 1993.** Effects of a native forest pathogen, *Phellinus weirii*, on Douglas-fir forest composition in western Oregon. *Canadian Journal of Forest Research*. 23: 2473–2480.
- Ingham, E.R.; Molina, R. 1991.** Interactions among mycorrhizal fungi, rhizosphere organisms, and plants. In: Barbosa, P.; Krischik, V.A.; Jones, C.G., eds. *Microbial mediation of plant-herbivore interactions*. Hoboken, NJ: John Wiley & Sons: 169–197.
- Jacobs, K.M.; Luoma, D.L. 2008.** Small mammal mycophagy response to variations in green-tree retention. *Journal of Wildlife Management*. 72(8): 1747–1755.
- Jonsson, L.; Dahlberg, A.; Wilsson, M.-C.; Zackrisson, O.; Kårén, O. 1999.** Ectomycorrhizal fungal communities in late-successional Swedish boreal forests, and their composition following wildfire. *Molecular Ecology*. 8: 205–215.
- Juutilainen, K.; Halme, P.; Kotiranta, H.; Mönkkönen M. 2011.** Size matters in studies of dead wood and wood-inhabiting fungi. *Fungal Ecology*. 4(5): 342–349.
- Jusino, M.A.; Lindner, D.L.; Banik, M.T.; Rose, K.R.; Walters, J.R. 2016.** Experimental evidence of a symbiosis between red-cockaded woodpeckers and fungi. *Proceedings of the Royal Society B: Biological Sciences*. 283 (1827). doi:10.1098/rspb.2016.0106.
- Kahl, T.; Arnstadt, T.; Baber, K.; Bässler, C.; Bauhus, J.; Borken, W.; Buscot, F.; Floren, A.; Heibl, C.; Hessenmöller, D.; Hofrichter, M.; Hoppe, B.; Kellner, H.; Krüger, D.; Linsenmair, K.E.; Matzner, E.; Otto, P.; Purahong, W.; Seilwinder, C.; Schulze, E.-D.; Wende, B.; Weisser, W.W.; Gossner, M. 2017.** Wood decay rates of 13 temperate tree species in relation to wood properties, enzyme activities and organismic diversities. *Forest Ecology and Management*. 391: 86–95.
- Kropp, B.R. 1982.** Rotten wood as mycorrhizal inoculum for containerized western hemlock. *Canadian Journal of Forest Research*. 12: 428–431.
- Kruys, N.; Fries, C.; Jonsson, B.G.; Lämås, T.; Ståhl, G. 1999.** Wood-inhabiting cryptogams on dead Norway spruce (*Picea abies*) trees in managed Swedish boreal forests. *Canadian Journal of Forest Research*. 29: 178–186.

- Kruys, N.; Jonsson, B.G. 1999.** Fine woody debris is important for species richness on logs in managed boreal spruce forests of northern Sweden. *Canadian Journal of Forest Research*. 29: 1295–1299.
- Kucuker, D.M.; Baskent, E.Z. 2017.** Impact of forest management intensity on mushroom occurrence and yield with a simulation-based decision support system. *Forest Ecology and Management*. 389: 240–248.
- Lassauce, A.; Paillet, Y.; Jactel, H.; Bouget, C. 2011.** Deadwood as a surrogate for forest biodiversity: meta-analysis of correlations between deadwood volume and species richness of saproxylic organisms. *Ecological Indicators*. 11(5): 1027–1039.
- Lehmkuhl, J.F.; Gould, L.E.; Cazares, E.; Hosford, D.R. 2004.** Truffle abundance and mycophagy by northern flying squirrels in eastern Washington forests. *Forest Ecology and Management*. 200 (1–3): 49–65.
- Lehmkuhl, J.F.; Kennedy, M.; Ford, E.D.; Singleton, P.H.; Gaines, W.L.; Lind, R.L. 2007.** Seeing the forest for the fuel: integrating ecological values and fuels management. *Forest Ecology and Management*. 246: 73–80.
- Löhmus, A. 2011.** Silviculture as a disturbance regime: the effects of clear-cutting, planting and thinning on polypore communities in mixed forests. *Journal of Forestry Research*. 16(3): 194–202.
- Luoma, D.L. 2001.** Monitoring of fungal diversity at the Siskiyou Integrated Research Site with special reference to the Survey and Manage species *Arcangeliella camphorata* (Singer & Smith) Pegler & Young. Research report. On file with: U.S. Department of Agriculture, Forest Service, Siskiyou National Forest, Chetco Ranger District, 539 Chetco Ave., Brookings, OR 97415.
- Luoma, D.L.; Trappe, J.M.; Claridge, A.W.; Jacobs, K.M.; Cazares, E. 2003.** Relationships among fungi and small mammals in forested ecosystems. In: Zabel, C.J.; Anthony, R.G., eds. *Mammal community dynamics: management and conservation in the coniferous forests of western North America*. Cambridge, United Kingdom: Cambridge University Press: 343–373.
- Manning, T.; Maguire, C.C.; Jacobs, K.M.; Luoma D.L. 2003.** Additional habitat, diet and range information for the white-footed vole (*Arborimus albipes*). *American Midland Naturalist*. 150(1): 115–122.

- Marcot, B.G. 2002.** An ecological functional basis for managing decaying wood for wildlife. In: Laudenslayer, W.F., Jr.; Shea, P.J.; Valentine, B.E.; Weatherspoon; C.P. Lisle, T.E., eds. Proceedings of the symposium on the ecology and management of dead wood in western forests. Gen. Tech. Rep. PSW-GTR-181. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station: 895–910.
- Marcot, B.G.; Pope, K.L.; Slauson, K.; Welsh, H.H.; Wheeler, C.A.; Reilly, M.J.; Zielinski, W.J. [N.d.]** Other species and biodiversity of older forests. Manuscript in preparation. On file with: Bruce G. Marcot, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, 620 SW Main St., Suite 400, Portland OR 97208.
- Maser, C.; Anderson, R.G.; Cromack, K., Jr.; Williams, J.T.; Martin, R.E. 1979.** Dead and down woody material. In: Thomas, J.W., ed. Wildlife habitats in managed forests: the Blue Mountains of Oregon and Washington. Agric. Handb. 553. Washington, DC: U.S. Department of Agriculture, Forest Service: 78–95.
- Maser, C.; Maser, Z. 1988.** Interactions among squirrels, mycorrhizal fungi, and coniferous forests in Oregon. *Great Basin Naturalist*. 48: 358–369.
- Maser, C.; Trappe, J.M.; Nussbaum, R.A. 1978.** Fungal-small mammal interrelationships with emphasis on Oregon coniferous forests. *Ecology*. 59: 799–809.
- Mellen, K.; Marcot, B.G.; Ohmann, J.L.; Waddell, K.L.; Willhite, E.A.; Hostetler, B.B.; Livingston, S.A.; Ogden, C. 2002.** DecAID: a decaying wood advisory model for Oregon and Washington. In: Laudenslayer, W.F., Jr.; Shea, P.J.; Valentine, B.E.; Weatherspoon; C.P.; Lisle, T.E., eds. Proceedings of the symposium on the ecology and management of dead wood in western forests. Gen. Tech. Rep. PSW-GTR-181. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station: 527–533.
- Molina, R. 2008.** Protecting rare, little known, old-growth forest-associated fungi in the Pacific Northwest USA: a case study in fungal conservation. *Mycological Research*. 112: 613–638.
- Molina, R.; Castellano, M.; O’Dell, T.; Smith, J.; Pilz, D.; Dreisbach, T.; Dunham S. 2001.** Conservation and management of forest fungi in the Pacific Northwestern United States: an integrated ecosystem approach. In: Moore, D.; Nauta, M.M.; Rotheroe, M., eds. *Fungal conservation: issues and solutions*. Cambridge, United Kingdom: Cambridge University Press: 19–63.

- Molina, R.; O'Dell, T.; Luoma, D.; Amaranthus, M.; Castellano, M.; Russell, K. 1993.** Biology, ecology, and social aspects of wild edible mushrooms in the forests of the Pacific Northwest: a preface to managing commercial harvest. Gen. Tech. Rep. PNW-GTR-309. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 42 p.
- Muller, M.M.; Varama, M.; Heinonen, J.; Hallaksela, A.M. 2002.** Influence of insects on the diversity of fungi in decaying spruce wood in managed and natural forests. *Forest Ecology and Management*. 166(1–3): 165–181.
- O'Dell, T.E.; Castellano, M.A.; Trappe, J.M. 1993.** Biology and application of ectomycorrhizal fungi. In: Metting, F.B., ed. *Soil microbial ecology: applications in agricultural and environmental management*. New York: Marcel Dekker: 379–416.
- O'Dell, T.E.; Luoma, D.L.; Molina, R.J. 1992.** Ectomycorrhizal fungal communities in young, managed and old growth Douglas-fir stands. *Northwest Environmental Journal*. 8: 166–168.
- O'Dell, T.E.; Smith, J.E.; Castellano, M.; Luoma, D. 1996.** Diversity and conservation of forest fungi. In: Pilz, D.; Molina, R., eds. *Managing forest ecosystems to conserve fungus diversity and sustain wild mushroom harvests*. Gen. Tech. Rep. PNW-GTR-371. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 5–18.
- O'Hanlon-Manners, D.L.; Kotanen, P.M. 2004.** Logs as refuges from fungal pathogens for seeds of eastern hemlock (*Tsuga canadensis*). *Ecology*. 85(1): 284–289.
- O'Neill, T.A.; Johnson, D.H.; Barrett, C.; Trevithick, M.; Bettinger, K.A.; Kiilsgaard, C.; Vander Heyden, M.; Greda, E.L.; Stinson, D.; Marcot, B.G.; Doran, P.J.; Tank, S.; Wunder, L. 2001.** Matrixes for wildlife-habitat relationships in Oregon and Washington. In: Johnson, D.H.; O'Neill, T.A., eds. *Wildlife-habitat relationships in Oregon and Washington*. [CD-ROM]. Corvallis OR: Oregon State University Press.
- Ohlson, M.; Söderström, L.; Hörnberg, G.; Zackrisson, O.; Hermansson, J. 1997.** Habitat qualities versus long-term continuity as determinants of biodiversity in boreal old-growth swamp forests. *Biological Conservation*. 81: 221–231.

- Osono, T. 2015.** Decomposing ability of diverse litter-decomposer macrofungi in subtropical, temperate, and subalpine forests. *Journal of Forest Research*. 20(2): 272–280.
- Paillet, Y.; Archaux, F.; Boulanger, V.; Debaive, N.; Fuhr, M.; Gilg, O.; Gosselin, F.; Guilbert, E. 2017.** Snags and large trees drive higher tree microhabitat densities in strict forest reserves. *Forest Ecology and Management*. 389: 176–186.
- Parladé, J.; Martínez-Peña, F.; Pera, J. 2017.** Effects of forest management and climatic variables on the mycelium dynamics and sporocarp production of the ectomycorrhizal fungus *Boletus edulis*. *Forest Ecology and Management*. 390: 73–79.
- Pilz, D.; Brodie, F.D.; Alexander, S.; Molina, R. 1998.** Relative value of chanterelles and timber as commercial forest products. *AMBIO Special Report No. 9*: 14–16.
- Pilz, D.P.; Perry, D.A. 1984.** Impact of clearcutting and slash burning on ectomycorrhizal associations of Douglas-fir seedlings. *Canadian Journal of Forest Research*. 14: 94–100.
- Rizzo, D.M.; Garbelotto, M. 2003.** Sudden oak death: endangering California and Oregon forest ecosystems. *Frontiers in Ecology and the Environment*. 1(5): 197–204.
- Rydin, H.; Kiekmann, M.; Hallingbäck, T. 1997.** Biological characteristics, habitat associations, and distribution of macrofungi in Sweden. *Conservation Biology*. 11: 628–640.
- Seibold, S.; Bäessler, C.; Brandl, R.; Gossner, M.M.; Thorn, S.; Ulyshen, M.D.; Müller, J. 2015.** Experimental studies of dead-wood biodiversity—a review identifying global gaps in knowledge. *Biological Conservation*. 191: 139–149.
- Schlosser, W.E.; Blatner K.A. 1995.** The wild edible mushroom industry of Washington, Oregon and Idaho: a 1992 survey. *Journal of Forestry*. 93: 31–36.
- Schowalter, T.D. 2000.** *Insect ecology: an ecosystem approach*. San Diego, CA: Academic Press. 483 p.
- Smith, J. E.; Kluber, L.A.; Jennings, T.N.; McKay, D.; Brenner, G.; Sulzman E.W. 2017.** Does the presence of large down wood at the time of a forest fire impact soil recovery? *Forest Ecology and Management*. 391: 52–62.

- Smith, J.E.; Molina, R.; Huso, M.M.P.; Larsen M.J. 2000.** Occurrence of *Piloderma fallax* in young, rotation-age, and old-growth stands of Douglas-fir (*Pseudotsuga menziesii*) in the Cascade Range of Oregon, U.S.A. *Canadian Journal of Botany*. 78: 995–1001.
- Smith, M.L.; Bruhn, J.N.; Anderson, J.B. 1992.** The fungus *Armillaria bulbosa* is among the largest and oldest organisms. *Nature*. 356: 428–431.
- Smith, S.E.; Read, D.J. 1997.** Mycorrhizal symbiosis. 2nd ed. San Diego, CA: Academic Press. 605 p.
- Spies, T.A.; Cline, S.P. 1988.** Coarse woody debris in forests and plantations of coastal Oregon. In: Mater, C.; Tarrant, R.; Trappe, J.M.; Franklin, J.F., eds. From the forest to the sea: a story of fallen trees. Gen. Tech. Rep. PNW-GTR-229. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 5–24. Chapter 1.
- Stendell, E.R.; Horton, T.R.; Bruns, T.D. 1999.** Early effects of prescribed fire on the structure of the ectomycorrhizal fungus community in a Sierra Nevada ponderosa pine forest. *Mycological Research*. 103: 1353–1359.
- Sverdrup-Thygeson, A.; Gustafsson, L.; Kouki, J. 2014.** Spatial and temporal scales relevant for conservation of dead-wood associated species: current status and perspectives. *Biodiversity and Conservation*. 23: 513–535.
- Swift, M.J. 1982.** Basidiomycetes as components of forest ecosystems. In: Frankland, J.C.; Hedger, J.N.; Swift, M.J., eds. Decomposer basidiomycetes: their biology and ecology. Cambridge, United Kingdom: Cambridge University Press: 307–338.
- Trofymow, J.A.; Addison, J.; Blackwell, B.A.; He, F.; Preston, C.A.; Marshall, V.G. 2003.** Attributes and indicators of old-growth and successional Douglas-fir forests on Vancouver Island. *Environmental Review*. 11 (Suppl. 1): S187–S204.
- Tyler, G. 1992.** Tree species affinity of decomposer and ectomycorrhizal macro-fungi in beech (*Fagus sylvatica* L.), oak (*Quercus robur* L.) and hornbeam (*Carpinus betulus* L.) forests. *Forest Ecology and Management*. 47: 269–284.
- Ulyshen, M.D. 2016.** Wood decomposition as influenced by invertebrates. *Biological Reviews*. 91: 70–85.

- van der Heijden, M.G.A.; Klironomos, J.N.; Ursic, M.; Moutoglis, P.; Streitwold-Engel, R.; Boller, T.; Wiemken, A.; Sanders, I.R. 1998.** Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. *Nature*. 396: 69–72.
- van der Wal, A.; Ottosson, E.; de Boer, W. 2015.** Neglected role of fungal community composition in explaining variation in wood decay rates. *Ecology*. 96(1): 124–133.
- Wästerlund, I.; Ingelög, T. 1981.** Fruit body production of larger fungi in some young Swedish forests with special reference to logging waste. *Forest Ecology and Management*. 3: 269–294.
- Weslien, J.; Djupström, L.B.; Schroeder, M.; Widenfalk, O. 2011.** Long-term priority effects among insects and fungi colonizing decaying wood. *Journal of Animal Ecology*. 80(6): 1155–1162.
- White, P.J.T.; McGill, B.J.; Lechowicz, M.J. 2012.** Detecting changes in forest floor habitat after canopy disturbance. *Ecological Research*. 27(2): 397–406.
- White, R.E. 1983.** A field guide to the beetles of North America. Boston, MA: Houghton Mifflin. 368 p.

Pacific Northwest Research Station

Website	http://www.fs.fed.us/pnw/
Telephone	(503) 808–2592
Publication requests	(503) 808–2138
FAX	(503) 808–2130
E-mail	pnw_pnwpubs@fs.fed.us
Mailing address	Publications Distribution Pacific Northwest Research Station P.O. Box 3890 Portland, OR 97208–3890



Federal Recycling Program
Printed on Recycled Paper

U.S. Department of Agriculture
Pacific Northwest Research Station
1220 SW 3rd Ave.
P.O. Box 3890
Portland, OR 97208-3890

Official Business
Penalty for Private Use, \$300